ONCE-WEEKLY
wegovy®
semaglutide injection 2.4 mg

Wegovy® is an injectable prescription medicine to treat adults with metabolic dysfunction-associated steatohepatitis (MASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver, along with a reduced calorie diet and increased physical activity.

Talking MASH with your health care professional

If you have been diagnosed with MASH with liver scarring, use this guide to help prepare for and track your conversations with your health care professional as you explore treatment.

Print this guide, write down what's important to you, and bring it to your next appointment.

When discussing a new medication you've been prescribed, you might have some questions. Your health care professional is your partner in creating a personalized treatment plan and is the best source of information on what treatment might be right for you. Once you decide on a treatment together, be sure to work with them on a plan.

Ask your health care professional these questions and use the space below to note their thoughts and recommendations.

1. I have MASH with liver scarring. Do I need any other tests or regular check-ups for my treatment plan?

2. How can medication be helpful for improving my MASH?

3. If Wegovy® is right for me, when and how should I take it? Is there anything else I should know?

4. What side effects should I watch out for while taking Wegovy®?

What is Wegovy®?

Wegovy® (semaglutide) injection 2.4 mg is an injectable prescription medicine used with a reduced calorie diet and increased physical activity to treat adults with metabolic dysfunction-associated steatohepatitis (MASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver.

This indication is approved based on improvement of MASH and liver scarring. There is an ongoing study to confirm the clinical benefit of Wegovy® in adults with MASH.

Wegovy® contains semaglutide and should not be used with other semaglutide-containing products or other GLP-1 receptor agonist medicines.

It is not known if Wegovy® is safe and effective for the treatment of MASH in people under 18 years of age.

Important Safety Information

What is the most important information I should know about Wegovy®?

Wegovy® may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, Wegovy® and medicines that work like Wegovy® caused thyroid tumors, including thyroid cancer. It is not known if Wegovy® will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people
- Do not use Wegovy® if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC) or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)

Please see additional Important Safety Information throughout. Please click here for Prescribing Information, including Medication Guide.

Do not use Wegovy® if:

- you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC) or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)
- you have had a serious allergic reaction to semaglutide or any of the ingredients in Wegovy®. See **"What are the possible side effects of Wegovy®?"** for symptoms of a serious allergic reaction

Wegovy® FlexTouch®: Do not share your Wegovy® FlexTouch® pen with other people, even if the needle has been changed. You may give other people a serious infection or get a serious infection from them.

Before using Wegovy®, tell your healthcare provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys
- have type 2 diabetes and a history of diabetic retinopathy
- are scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation)
- are pregnant or plan to become pregnant. Wegovy® may harm your unborn baby. You should stop using Wegovy® 2 months before you plan to become pregnant
- are breastfeeding or plan to breastfeed. It is not known if Wegovy® when received through an injection passes into your breast milk

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Wegovy® may affect the way some medicines work and some medicines may affect the way Wegovy® works. Tell your healthcare provider if you are taking other medicines to treat diabetes, including sulfonylureas or insulin. Wegovy® slows stomach emptying and can affect medicines that need to pass through the stomach quickly.

Tips for a productive conversation
with your health care professional

Write down any other questions or concerns.

Appointments can go by quickly, so having a list helps your health care professional focus on what's important to you.

Share any symptoms.

Let your health care professional know what you're noticing, thinking, and feeling so they can give you the best care.

Take notes.

Reviewing your notes later can help you remember important instructions or information your health care professional provided.

Write down any additional questions you may have here:

Important Safety Information (cont'd)

What are the possible side effects of Wegovy®?

Wegovy® may cause serious side effects, including:

- **inflammation of your pancreas (pancreatitis).** Stop using Wegovy® and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without nausea or vomiting. Sometimes you may feel the pain from your abdomen to your back
- **gallbladder problems.** Wegovy® may cause gallbladder problems, including gallstones. Some gallstones may need surgery. Call your healthcare provider if you have symptoms, such as pain in your upper stomach (abdomen), fever, yellowing of the skin or eyes (jaundice), or clay-colored stools
- **increased risk of low blood sugar (hypoglycemia), especially those who also take medicines for diabetes such as insulin or sulfonylureas.** This can be a serious side effect. Talk to your healthcare provider about how to recognize and treat low blood sugar and check your blood sugar before you start and while you take Wegovy®. Signs and symptoms of low blood sugar may include dizziness or light-headedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion or drowsiness, shakiness, weakness, headache, fast heartbeat, or feeling jittery
- **dehydration leading to kidney problems.** Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration. Tell your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away
- **severe stomach problems.** Stomach problems, sometimes severe, have been reported in people who use Wegovy®. Tell your healthcare provider if you have stomach problems that are severe or will not go away
- **serious allergic reactions.** Stop using Wegovy® and get medical help right away, if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat; problems breathing or swallowing; severe rash or itching; fainting or feeling dizzy; or very rapid heartbeat
- **change in vision in people with type 2 diabetes.** Tell your healthcare provider if you have changes in vision during treatment with Wegovy®
- **increased heart rate.** Wegovy® can increase your heart rate while you are at rest. Tell your healthcare provider if you feel your heart racing or pounding in your chest and it lasts for several minutes
- **food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).** Wegovy® may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking Wegovy® before you are scheduled to have surgery or other procedures

The most common side effects of Wegovy® may include: nausea, diarrhea, vomiting, constipation, stomach (abdomen) pain, changes in skin sensations, headache, tiredness (fatigue), upset stomach, dizziness, feeling bloated, belching, low blood sugar in people with type 2 diabetes, gas, stomach flu, heartburn, and hair loss.

Please see additional Important Safety Information on the previous page, and click here for Prescribing Information and Medication Guide.

Wegovy® is a prescription medication. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit **www.fda.gov/medwatch,** or call **1-800-FDA-1088.**

FDA, US Food and Drug Administration.

ONCE-WEEKLY
wegovy®
semaglutide injection **2.4 mg**

Wegovy® and FlexTouch® are registered trademarks of Novo Nordisk A/S.
Novo Nordisk is a registered trademark of Novo Nordisk A/S.
© 2026 Novo Nordisk All rights reserved.
US26SN00133 August 2026

novo nordisk®